

Construction of Doubly Periodic Solutions via the Poincaré–Lindstedt Method in the case of Massless φ^4 Theory

Oleg A. Khrustalev

*Bogoliubov Institute for Theoretical Problems of Microphysics,
Moscow State University;
Department of Physics,
Moscow State University,
Vorobievsky Gory, Moscow, 119899, RUSSIA*
E-mail: khrust@sunny.bog.msu.su

and

Sergey Yu. Vernov

*Skobeltsyn Institute for Nuclear Physics,
Moscow State University,
Vorobievsky Gory, Moscow, 119899, RUSSIA*
E-mail: svernov@theory.npi.msu.su

Abstract

Doubly periodic (periodic both in time and in space) solutions for the Lagrange–Euler equation of the $(1 + 1)$ –dimensional scalar φ^4 theory are considered. The nonlinear term is assumed to be small, and the Poincaré–Lindstedt method is used to find asymptotic solutions in the standing wave form. The principal resonance problem, which arises for zero mass, is solved if the leading-order term is taken in the form of a Jacobi elliptic function. It has been proved that the choice of elliptic cosine with fixed value of module k ($k \approx 0.451075598811$) as the leading-order term puts the principal resonance to zero and allows us to construct (with accuracy to third order of small parameter) the asymptotic solution in the standing wave form. To obtain this leading-order term the computer algebra system **REDUCE** has been used. We have appended the **REDUCE** program to this paper.

1 Introduction

1.1 Periodic solutions of nonlinear equations

Recently, periodic solutions of the nonlinear wave equation:

$$\frac{\partial^2 \varphi(x, t)}{\partial x^2} - \frac{\partial^2 \varphi(x, t)}{\partial t^2} - g(\varphi) = 0, \quad (1)$$

where $g(\varphi)$ denotes a continuous function on $\mathbf{R}$ such that $g(0) = 0$, are discussed intensively. In the 1970s and 1980s existence theorems for a wide class of continuous functions $g(\varphi)$ satisfying definite boundary conditions in x have been proved.

J. M. Coron [1] has proved that the following problem:

$$\begin{cases} \frac{\partial^2 \varphi(x, t)}{\partial x^2} - \frac{\partial^2 \varphi(x, t)}{\partial t^2} - g(\varphi) = 0, & x, t \in \mathbf{R}, \quad g(0) = 0, \\ \lim_{|x| \rightarrow \infty} \varphi(x, t) = 0, & x, t \in \mathbf{R}, \\ \varphi(x, t + T) = \varphi(x, t), & x, t, T \in \mathbf{R}, \end{cases} \quad (2)$$

has a T -periodic (in time) solutions only if $g'(0) > \left(\frac{2\pi}{T}\right)^2$.

In particular, if $g(\varphi) = \sin(\varphi)$, there are no T -periodic solutions when $T < 2\pi$ (for any period $T > 2\pi$ an explicit solution is known, see e.g. G. Lamb [2]). If $g(\varphi) = -\sin(\varphi)$ or $g(\varphi) = \pm\varphi^3$, then there is no nontrivial solution of problem (2) (for any period T). To obtain periodic solutions of massless φ^4 theory one has to change the boundary conditions.

Generalizing the Rabinowitz theorem [3], H. Brezis, J. M. Coron and L. Nirenberg [4] have proved that for each T , which is a rational multiple of π , the following problem:

$$\begin{cases} \frac{\partial^2 \varphi(x, t)}{\partial x^2} - \frac{\partial^2 \varphi(x, t)}{\partial t^2} - g(\varphi) = 0, & x, t \in \mathbf{R}, \quad g(0) = 0, \\ \varphi(0, t) = \varphi(\pi, t) = 0, & t \in \mathbf{R}, \\ \varphi(x, t + T) = \varphi(x, t), & x, t, T \in \mathbf{R}, \end{cases} \quad (2')$$

has a nontrivial weak T -periodic solution if $g(\varphi)$ is a continuous nondecreasing function, satisfying the following conditions:

$$\lim_{|\varphi| \rightarrow \infty} \frac{g(\varphi)}{\varphi} = \infty,$$

$$\exists \alpha, \beta > 0 \quad \text{such that} \quad \forall \varphi \in \mathbf{R} : \quad \frac{1}{2} \varphi \cdot g(\varphi) - \int_0^\varphi g(\psi) d\psi \geq \beta |g(\varphi)| - \alpha.$$

By a *weak solution* we mean such function φ that

$$\int_0^T \int_0^\pi (\varphi(v_{tt} - v_{xx}) + g(\varphi)v) dt dx = 0$$

for all $v \in C^2([0, \pi] \times \mathbf{R})$ satisfying the boundary and periodicity conditions. A simplified proof of the Rabinowitz theorem for the case

$$g(\varphi) = |\varphi|^n \cdot \varphi,$$

where $n > 0$, was given in [5].

1.2 Two kinds of doubly periodic solutions

Our investigation is dedicated to the construction of doubly periodic classical fields in the $(1+1)$ -dimensional φ^4 theory. We study the model of an isolated real scalar field $\varphi(x, t)$, described by the Lagrangian density:

$$\mathcal{L}(\varphi) = \frac{1}{2} \left(\varphi_{,t}^2(x, t) - \varphi_{,x}^2(x, t) - M^2 \varphi^2(x, t) - \frac{\varepsilon}{2} \varphi^4(x, t) \right).$$

The corresponding Lagrange–Euler equation is:

$$\frac{\partial^2 \varphi(x, t)}{\partial x^2} - \frac{\partial^2 \varphi(x, t)}{\partial t^2} - M^2 \varphi(x, t) - \varepsilon \varphi^3(x, t) = 0. \quad (3)$$

There are two classes of doubly periodic solutions for this equation. If we seek fields in the traveling wave form:

$$\varphi(x, t) = \varphi(x - vt),$$

where v is the velocity of the wave motion, then equation (3) reduces to the Duffing's equation [6]. As is known [7], periodic solutions for this equation are Jacobi elliptic functions. Such solutions are well known for both $(1+1)$ -dimensional [8] and $(3+1)$ -dimensional [9] variants of φ^4 theory. They are used in the Yang–Mills theory [10].

The second class of doubly periodic solutions consists of functions in the standing wave form:

$$\varphi(x, t) \equiv \sum_{n=1}^{\infty} \sum_{j=1}^{\infty} \mathcal{C}_{nj} \sin(n(x - x_0)) \sin(j \omega (t - t_0)), \quad (4)$$

where x_0 and t_0 are constants determined by boundary and initial conditions. Equation (3) is a translation-invariant, so we can restrict our consideration to the case of zero x_0 and t_0 without loss of generality. The choice $x_0 = 0$ corresponds to the boundary conditions of problem (2'). The period of $\varphi(x, t)$ in x is taken to be 2π and we are interesting in finding the period in t .

Exact standing wave solutions are not known. Approximate solutions for equation (3) with $\varepsilon = 1$ and $M = 1$ have been found under the assumption that all highest harmonics are zeros: $\forall n, j > N : \mathcal{C}_{nj} = 0$, where N is a large number. Numerical calculations have been made [11] for several values of N . The obtained solutions are very close to each other, but convergence of the sequence of these solutions as N tends to infinity cannot be proved in numerical framework.

2 Asymptotic solutions in the standing wave form

Let us consider periodic solutions for equation (3), provided that ε is small or, equivalently, that these solutions have small amplitudes.

An asymptotic expansion, containing only bounded functions, is called a *uniform expansion*. The possibility to obtain a uniform expansion, using standard asymptotic methods, for example, Poincaré–Lindstedt [12] or Krylov–Bogoliubov [13] methods, depends on values of the frequencies in the zero approximation (for a review of such methods, see [14, 15]). If $\varepsilon = 0$, then equation (3) is linear and has periodic solutions of the form (4)

with frequencies in time $\Omega_j = \sqrt{j^2 + M^2}$, where $j \in \mathbb{N}$. There are two fundamentally different cases.

If

$$\forall i, j \in \mathbb{N} \quad \nexists k, n \in \mathbb{N} \quad \text{such that} \quad \frac{\Omega_i}{\Omega_j} = \frac{k}{n},$$

then it is a nonresonance case and a uniform expansion can be found with any desired accuracy using the standard asymptotic methods, for instance, the Poincaré–Lindstedt method. All secular terms can be eliminated with the proper choice of frequency.

The resonance case, when there exist two frequencies Ω_j and Ω_i , whose ratio is rational, is more difficult. The Krylov–Bogoliubov method and the standard variant of the Poincaré–Lindstedt method can be used to find periodic solutions only to a few leading orders in ε .

The important example of resonance case is the massless φ^4 theory when $\Omega_j = j$ and all relations of frequencies are rational, which is the principal resonance case. Using standard asymptotic methods, one cannot construct a periodic solution even to the first order in ε . It is possible to transform the differential equations to a system of nonlinear algebraic equations in Fourier coefficients and frequencies using the Poincaré–Dulac’s normal form method. The algorithm of this procedure was constructed [16, 17] and realized in the software for symbolic and algebraic computation **REDUCE** [18–20] (the system **NORT** [21, 22]). But an algorithm to solve the resulting algebraic system has yet to be created.

3 The standing wave solutions for the massless φ^4 theory

3.1 Construction of asymptotic solutions via the Poincaré–Lindstedt method

The purpose of this article is the construction of standing wave solutions of equation (3) with $M = 0$, using asymptotic methods when $\varepsilon \ll 1$. We use the Poincaré–Lindstedt method: introduce the new time $\tilde{t} \equiv \omega t$ and look for a doubly periodic function $\varphi(x, \tilde{t})$ and a frequency (in time) ω in the form of power series in ε :

$$\begin{aligned} \varphi(x, \tilde{t}, \varepsilon) &\equiv \sum_{n=0}^{\infty} \varphi_n(x, \tilde{t}) \varepsilon^n, \\ \omega(\varepsilon) &\equiv 1 + \sum_{n=1}^{\infty} \omega_n \varepsilon^n. \end{aligned}$$

Expanding the Lagrange–Euler equation in a power series in ε , we obtain a series of equations. The first two equations of this series are:

- to zero order in ε the equation for φ_0 is:

$$\frac{\partial^2 \varphi_0(x, \tilde{t})}{\partial x^2} - \frac{\partial^2 \varphi_0(x, \tilde{t})}{\partial \tilde{t}^2} = 0, \tag{5}$$

- to first order in ε the equation in φ_0 , φ_1 and ω_1 is:

$$\frac{\partial^2 \varphi_1(x, \tilde{t})}{\partial x^2} - \frac{\partial^2 \varphi_1(x, \tilde{t})}{\partial \tilde{t}^2} = 2\omega_1 \frac{\partial^2 \varphi_0(x, \tilde{t})}{\partial \tilde{t}^2} + \varphi_0^3(x, \tilde{t}). \quad (6)$$

Equation (5) has many periodic solutions. If we select $\varphi_0(x, \tilde{t}) = \sin(x) \sin(\tilde{t})$, then the second equation has no periodic solution, because the frequency of the external force $\sin(3x) \sin(3\tilde{t})$ is equal to the frequency of its own oscillations and it is impossible to put this resonance harmonic to zero by selecting only ω_1 . In the massless case to obtain a uniform expansion, we must find another doubly periodic solution to equation (5). The correct selection of not only the frequency $\omega(\varepsilon)$, but also the function $\varphi_0(x, \tilde{t})$, allows us to construct the next to leading order term in standing wave form.

3.2 The condition of existence of periodic solutions

The general solution, in the standing wave form (4), for equation (5) is the function

$$\varphi_0(x, \tilde{t}) = \sum_{n=1}^{\infty} a_n \sin(nx) \sin(n\tilde{t})$$

with arbitrary a_n . We have to find coefficients a_n so that the function $\varphi_1(x, \tilde{t})$ is a periodic solution for equation (6). If we select $\varphi_1(x, \tilde{t})$ as a double sum:

$$\varphi_1(x, \tilde{t}) \equiv \sum_{n=1}^{\infty} \sum_{j=1}^{\infty} b_{nj} \sin(nx) \sin(j\tilde{t})$$

with arbitrary b_{nj} , then equation (6) can be presented in the form of Fourier series:

$$\frac{\partial^2 \varphi_1(x, \tilde{t})}{\partial x^2} - \frac{\partial^2 \varphi_1(x, \tilde{t})}{\partial \tilde{t}^2} - 2\omega_1 \frac{\partial^2 \varphi_0(x, \tilde{t})}{\partial \tilde{t}^2} - \varphi_0^3(x, \tilde{t}) = \sum_{n=1}^{\infty} \sum_{j=1}^{\infty} R_{nj}(\mathbf{a}, \mathbf{b}) \sin(nx) \sin(j\tilde{t}) = 0$$

and is equivalent to the following infinite system of the algebraic equations in Fourier coefficients of the functions $\varphi_0(x, \tilde{t})$ and $\varphi_1(x, \tilde{t})$:

$$\forall n, j : R_{nj}(\mathbf{a}, \mathbf{b}) = 0. \quad (6')$$

This system has a subsystem of the equations in the Fourier coefficients of the function $\varphi_0(x, \tilde{t})$:

$$\forall j \in \mathbb{N} : R_{jj}(\mathbf{a}) = 0, \quad (7)$$

where

$$\begin{aligned} R_{jj} \equiv & 9a_j^3 + 3a_j^2 a_{3j} + a_j \left(6 \sum_{\substack{s=1 \\ s \neq j}}^{\infty} (2a_s^2 + a_s a_{2j+s}) + 3 \sum_{\substack{s=1 \\ s \neq j}}^{2j-1} a_s a_{2j-s} - 32j^2 \omega_1 \right) + \\ & + 3 \sum_{\substack{s=1 \\ s \neq j}}^{\infty} \sum_{\substack{p=1 \\ p \neq j}}^{\infty} a_s a_p a_{j+s+p} + 3 \sum_{\substack{s=1 \\ s \neq j}}^{\infty} \sum_{\substack{p=1 \\ p \neq j \\ p \neq 2j-s}}^{\infty} a_s a_p a_{s+p-j} + \sum_{s=1}^{j-2} \sum_{p=1}^{j-2} a_s a_p a_{j-s-p} \end{aligned}$$

If the system (7) is solved and all Fourier coefficients of the function $\varphi_0(x, \tilde{t})$ are known, then the system (6') transforms into a system of linear equations: $R_{nj}(\mathbf{b}) = 0$, which always has only one solution. Hence, we have obtained the necessary and sufficient condition of the existence of periodic solutions for equation (6): there exists a periodic function $\varphi_1(x, \tilde{t})$, satisfying equation (6), if and only if the Fourier coefficients of the function $\varphi_0(x, \tilde{t})$ satisfy system (7).

The coefficient a_1 is a parameter determining the oscillation amplitude. In fact, let $a_j = c_j a_1$ and $\omega_1 = c_\omega a_1^2$, then all polynomials $R_{jj}(\mathbf{a})$ are proportional to a_1^3 : $R_{jj}(\mathbf{a}) = a_1^3 R_{jj}(\mathbf{c})$ and, therefore, the coefficient a_1 can be selected arbitrarily.

3.3 The approximate solution of system (7)

System (7) is very difficult to solve. On the one hand, all R_{jj} are infinite series and the number of equations is infinite too. On the other hand, each equation of this system is nonlinear. We restrict ourselves to finding a particular solution. To simplify calculations, we assume that the function $\varphi_0(x, \tilde{t})$ contains only odd harmonics. Our goal is to find a real solution so we seek $c_j \in \mathbb{R}$.

To find an approximate solution we apply the Galerkin method: cut off higher diagonal harmonics $\forall j > N$: $c_{2j-1} = 0$ and seek an approximation for $\varphi_0(x, \tilde{t})$ in the following form

$$\varphi_0(x, \tilde{t}) = a_1 \left\{ \sum_{j=1}^N c_{2j-1} \sin((2j-1)x) \sin((2j-1)\tilde{t}) \right\}.$$

Neglecting the higher harmonics of the function $\varphi_0(x, \tilde{t})$, we obtain a finite system of nonlinear equations, with the number of equations three times as many as that of the variables. We solve the system of the N leading equations and substitute the obtained values in the other equations. In principle, any system of N nonlinear equations in N unknowns can be solved with a computer, using Buchberger algorithm [23, 24]. This algorithm has been realized in the standard procedures of the computer algebra system **REDUCE**: the **SOLVE** operator and the Gröebner basis package. This algorithm allows us to diagonalize this system of nonlinear equations, constructing an equivalent system, which consists of equation in only one variable, equation in two variables and so on. So, we can obtain solution of the system, solving only equations in one variable and substitute the result into the remaining equations.

The operating memory and other parameters of real computers are not enough to solve very difficult systems. For example, using computer with 128 Mbytes operating memory (RAM), we have solved the system, consisting of the N leading equations of the system (7) only for $N < 6$. On the other hand, even for these values of N the solution of system can be found only with finite accuracy (the exact solution can be irrational).

The **SOLVE** operator in **REDUCE** (in "on rounded" mode) rounds off numbers with accuracy $\delta = 10^{-11}$. We assume that the system of the N leading equations is solved if for all $j \leq N$ the inequalities $|R_{jj}(\mathbf{c})| < \delta$ are true. We also admit that the value of N is sufficient to solve system (7) if for all $j \in \mathbb{N}$ the inequalities $|R_{jj}(\mathbf{c})| < \delta$ are true.

To find a particular solution of the N leading equations for any $N < 50$ we have written a short program in **REDUCE** (see *Appendix*) and obtained that the minimal

sufficient value of N for $\delta = 10^{-11}$ is $N = 8$ and that the frequency correction is

$$\omega_1 = 0.28268003454 a_1^2.$$

We also have found numerical values of the fifteen leading Fourier coefficients of $\varphi_0(x, \tilde{t})$. The values of c_j are very close to the values of the corresponding terms of the following finite sequence:

$$\mathbf{d} = \{d_{2j-1} = \frac{f_{2j-1}}{f_1}, \text{ where } f_{2j-1} \equiv \frac{q^{j-1/2}}{1 + q^{2j-1}}; d_{2j} = 0; j < 23\},$$

with $q = 0.0142142623201$. It is easy to verify that substitution of this finite sequence gives $\forall j \in \mathbb{N} : |R_{jj}(\mathbf{d})| < 10^{-12}$.

In other words, the finite sequence $\mathbf{d}$ is an approximate solution of system (7). These numerical calculations help to find the analytical form of $\varphi_0(x, \tilde{t})$. The following table illustrates this interesting result:

j	c_j	$R_{jj}(\mathbf{c})$	d_j	$R_{jj}(\mathbf{d})$
1	1	0	1	0
3	$1.44162661711 \times 10^{-2}$	3.5×10^{-12}	$1.44162661711 \times 10^{-2}$	-8.0×10^{-14}
5	$2.04917177408 \times 10^{-4}$	2.1×10^{-12}	$2.04917177419 \times 10^{-4}$	-3.4×10^{-15}
7	$2.91274649724 \times 10^{-6}$	7.8×10^{-12}	$2.91274651543 \times 10^{-6}$	-9.7×10^{-17}
9	$4.14025418115 \times 10^{-8}$	8.3×10^{-13}	$4.14025430425 \times 10^{-8}$	-2.3×10^{-18}
11	$5.88506592014 \times 10^{-10}$	1.0×10^{-14}	$5.88506607528 \times 10^{-10}$	-4.9×10^{-20}
13	$8.36488192079 \times 10^{-12}$	4.6×10^{-13}	$8.36518729655 \times 10^{-12}$	-9.7×10^{-22}
15	$1.18901919266 \times 10^{-13}$	-7.8×10^{-22}	$1.1890496659 \times 10^{-13}$	-1.8×10^{-23}
17	0	4.4×10^{-12}	$1.69014638629 \times 10^{-15}$	-3.4×10^{-25}
19	0	7.3×10^{-14}	$2.40241840942 \times 10^{-17}$	-6.0×10^{-27}
21	0	1.1×10^{-15}	$3.41486054743 \times 10^{-19}$	-1.0×10^{-28}
23	0	1.7×10^{-17}	$4.85397236079 \times 10^{-21}$	-1.8×10^{-30}
25	0	2.4×10^{-19}	$6.89956364312 \times 10^{-23}$	-3.0×10^{-32}
27	0	3.3×10^{-21}	$9.8072207518 \times 10^{-25}$	-4.9×10^{-34}
29	0	4.2×10^{-23}	$1.39402408398 \times 10^{-26}$	-8.1×10^{-36}
31	0	5.2×10^{-25}	$1.98150240103 \times 10^{-28}$	-1.3×10^{-37}
33	0	5.7×10^{-27}	$2.81655949163 \times 10^{-30}$	-2.1×10^{-39}
35	0	6.1×10^{-29}	$4.00353154544 \times 10^{-32}$	-3.4×10^{-41}
37	0	6.2×10^{-31}	$5.69072475939 \times 10^{-34}$	-5.4×10^{-43}
39	0	5.9×10^{-33}	$8.08894545219 \times 10^{-36}$	-8.5×10^{-45}
41	0	5.0×10^{-35}	$1.14978392551 \times 10^{-37}$	-1.3×10^{-46}
43	0	3.6×10^{-37}	$1.63433303287 \times 10^{-39}$	-2.1×10^{-48}
45	0	1.7×10^{-39}	$2.32308384477 \times 10^{-41}$	-1.6×10^{-49}
47	0	0	0	1.4×10^{-50}

3.4 The exact solution of system (7)

The approximate solution of system (7) is found. The aim of this section is to find an exact solution of (7), that is to say to find the function $\varphi_0(x, t)$ in analytical form. For

arbitrary $q \in (0, 1)$ let us define the following sequence:

$$\mathbf{f} \stackrel{\text{def}}{=} \{ \forall n \in \mathbb{N} : f_{2n-1} = \frac{q^{n-1/2}}{1+q^{2n-1}}, f_{2n} = 0 \}.$$

The terms of the sequence $\mathbf{f}$ are proportional to Fourier coefficients of the Jacobi elliptic function cn (the elliptic cosine):

$$\text{cn}(z, k) = \frac{\gamma}{k} \sum_{n=1}^{\infty} f_{2n-1} \cos \left((2n-1) \frac{\gamma z}{4} \right), \quad \text{where } \gamma \equiv \frac{2\pi}{K}, \quad z \in \mathbb{R}. \quad (8)$$

Let us clarify the notation and point out some properties of the elliptic cosine [7]:

- Basic periods of the doubly periodic function $\text{cn}(z, k)$ are $4K(k)$ and $2K(k) + 2iK'(k)$, where $K(k)$ is a full elliptic integral, $K'(k) \equiv K(k')$ and $k' = \sqrt{1-k^2}$.
- The parameter q in the Fourier expansion can be expressed in terms of elliptic integrals: $q \equiv e^{-\pi \frac{K'}{K}}$.
- The Fourier-series expansion of the function $\text{cn}(z, k)$ does not include even harmonics. This expansion is valid in the following domain of the complex plane: $-K' < \Im m z < K'$, in particular, for $z \in \mathbb{R}$.
- If $z \in \mathbb{R}$ and $k \in (0, 1)$, then $\text{cn}(z, k) \in \mathbb{R}$.
- The function $\text{cn}(z, k)$ is a solution of the following differential equation:

$$\frac{d^2 \text{cn}(z, k)}{dz^2} = (2k^2 - 1) \text{cn}(z, k) - 2k^2 \text{cn}^3(z, k). \quad (9)$$

The latest property means that the infinite sequence of the Fourier coefficients for $\text{cn}(z, k)$ is a solution of some infinite system of nonlinear algebraic equations. Let us find this system. On the one hand, it is clear from (8) that the Fourier-series expansion for the function $\text{cn}^3(z, k)$ is:

$$\begin{aligned} \text{cn}^3(z, k) &= \frac{\gamma^3}{4k^3} \sum_{j=1}^{\infty} F_j^{(3)}(\mathbf{f}) \cos \left(j \frac{\gamma z}{4} \right), \quad \text{where } j = 1, 3, 5, \dots, +\infty ; \\ F_j^{(3)}(\mathbf{f}) &= 3f_j^3 + 3f_j^2 f_{3j} + f_j \left(6 \sum_{\substack{s=1 \\ s \neq j}}^{\infty} (f_s^2 + f_s f_{2j+s}) + 3 \sum_{\substack{s=1 \\ s \neq j}}^{2j-1} f_s f_{2j-s} \right) + \\ &+ 3 \sum_{\substack{s=1 \\ s \neq j}}^{\infty} \sum_{\substack{p=1 \\ p \neq j}}^{\infty} f_s f_p f_{j+s+p} + 3 \sum_{\substack{s=1 \\ s \neq j}}^{\infty} \sum_{\substack{p=1 \\ p \neq j \\ p \neq 2j-s}}^{\infty} f_s f_p f_{s+p-j} + \sum_{s=1}^{j-2} \sum_{p=1}^{j-2} f_s f_p f_{j-s-p} \end{aligned}$$

(in all sums we summarize over only odd numbers).

On the other hand, from the differential equation (9) it follows that $F_j^{(3)}(\mathbf{f})$ is proportional to f_j , with coefficients of proportionality depending on j :

$$\forall j \quad : \quad F_j^{(3)}(\mathbf{f}) = \left(\frac{2(2k^2 - 1)}{\gamma^2} + \frac{j^2}{8} \right) f_j. \quad (10)$$

Thus, the sequence $\mathbf{f}$ is a nonzero solution of system (10) at all $q \in (0, 1)$. The following lemma proves the existence of the selected value of q .

Lemma. *There exists such value of parameter $q \in (0, 1)$ that the sequence $\mathbf{f}$ becomes a real solution to system (7) with the real-valued parameter ω_1 .*

Proof. Inserting the sequence $\mathbf{f}$ into system (7) : $a_j = f_j$ and using system (10), we obtain:

$$\begin{aligned} \forall j \quad : \quad R_{jj}(\mathbf{f}) &= \left\{ F_j^{(3)}(\mathbf{f}) + f_j \left(6 \sum_{n=1}^{\infty} f_n^2 - 32j^2\omega_1 \right) \right\} = \\ &= f_j \left\{ 6 \sum_{n=1}^{\infty} f_n^2 + \frac{2(2k^2 - 1)}{\gamma^2} + j^2 \left(\frac{1}{8} - 32\omega_1 \right) \right\} = 0. \end{aligned}$$

System (7) has a nonzero solution if and only if

$$\begin{cases} \omega_1 = \frac{1}{256}, \\ \sum_{n=1}^{\infty} f_n^2 = \frac{(1 - 2k^2)}{3\gamma^2}. \end{cases}$$

We have obtained the value of ω_1 . The second equation of this system is equivalent to the following equation in parameter q :

$$3 \sum_{n=1}^{\infty} \left(\frac{q^{n-1/2}}{1 + q^{2n-1}} \right)^2 - \left(\frac{1}{4} + \sum_{n=1}^{\infty} \frac{q^n}{1 + q^{2n}} \right)^2 + 2 \left(\sum_{n=1}^{\infty} \frac{q^{n-1/2}}{1 + q^{2n-1}} \right)^2 = 0. \quad (11)$$

This equation has the following solution on interval (0,1):

$$q = 1.42142623201 \times 10^{-2} \pm 1 \times 10^{-13}.$$

Thus the lemma is proved.

We can now construct the desired leading-order approximation for the function $\varphi(x, \tilde{t})$:

$$\varphi_0(x, \tilde{t}) = A \left\{ \text{cn}(\alpha(x - \tilde{t}), k) - \text{cn}(\alpha(x + \tilde{t}), k) \right\}.$$

For arbitrary $k \in (0, 1)$ this function is a real solution of equation (6). If $\alpha = \frac{2K}{\pi}$, then the periods of $\varphi_0(x, \tilde{t})$ in x and in $\tilde{t}$ are equal to 2π . Using the Fourier-series expansion for the function $\text{cn}(z, k)$ (formula (8)), we obtain the following expansion for function $\varphi_0(x, \tilde{t})$:

$$\varphi_0(x, \tilde{t}) = 2A \frac{\gamma}{k} \sum_{n=1}^{\infty} f_{2n-1} \sin((2n-1)x) \sin((2n-1)\tilde{t}).$$

If $q = 1.42142623201 \times 10^{-2} \pm 1 \times 10^{-13}$, then q is a solution of equation (11) and the sequence $\mathbf{f}$ is a real solution of system (7). The middle value of q corresponds to $k = 0.451075598811$, $\gamma = 3.78191440007$ and $\alpha = 1.0576653982$. All equations in system (7) are homogeneous, hence for these values of parameters, the sequence of the Fourier coefficients of the function $\varphi_0(x, \tilde{t})$ also is a solution of system (7), with

$$\omega_1 = \frac{\gamma^2}{64k^2} A^2 = 1.0983600974 A^2.$$

Thus we have proved that the function

$$\varphi_0(x, \tilde{t}) = A \left\{ \text{cn}(\alpha(x - \tilde{t}), k) - \text{cn}(\alpha(x + \tilde{t}), k) \right\},$$

with $k = 0.451075598811$ and $\alpha = 1.0576653982$ is such standing wave solution of equation (5) that equation (6) has a periodic solution¹.

3.5 The first approximation

Now it is easy to find the periodic solution $\varphi_1(x, \tilde{t})$. Let us designate the Fourier coefficients of the function $\varphi_0^3(x, \tilde{t})$ as D_{nj} :

$$\varphi_0^3(x, \tilde{t}) \equiv \sum_{n=1}^{\infty} \sum_{j=1}^{\infty} D_{nj} \sin(nx) \sin(j\tilde{t}).$$

Equation (6) gives the following result (b_{nn} are arbitrary numbers):

$$\varphi_1(x, \tilde{t}) = \sum_{n=1}^{\infty} \sum_{\substack{j=1 \\ j \neq n}}^{\infty} \frac{D_{nj}}{j^2 - n^2} \sin(nx) \sin(j\tilde{t}) + \sum_{n=1}^{\infty} b_{nn} \sin(nx) \sin(n\tilde{t}).$$

It should be noted that the function $\varphi_1(x, \tilde{t})$ with arbitrary diagonal coefficients b_{nn} is a solution of equation (6) and that all off-diagonal coefficients of $\varphi_1(x, \tilde{t})$ are proportional to A^3 .

3.6 The second approximation

Let us consider the approximation to second order in ε :

$$\frac{\partial^2 \varphi_2(x, \tilde{t})}{\partial x^2} - \frac{\partial^2 \varphi_2(x, \tilde{t})}{\partial \tilde{t}^2} = 2\omega_1 \frac{\partial^2 \varphi_1(x, \tilde{t})}{\partial \tilde{t}^2} + (2\omega_2 + \omega_1^2) \frac{\partial^2 \varphi_0(x, \tilde{t})}{\partial \tilde{t}^2} + 3\varphi_1(x, \tilde{t})\varphi_0^2(x, \tilde{t}). \quad (12)$$

If all diagonal coefficients of $\varphi_1(x, \tilde{t})$ are zeros: $\forall n : b_{nn} = 0$, then $\forall j, n : b_{jn} = -b_{nj}$, and the function $\varphi_1(x, \tilde{t})\varphi_0^2(x, \tilde{t})$ has no diagonal harmonics. Hence, selecting $\omega_2 = -\frac{1}{2}\omega_1^2$, we obtain a periodic solution to equation (12):

¹This result has been published in our paper [25]

$$\varphi_2(x, \tilde{t}) \equiv \sum_{n=1}^{\infty} \sum_{\substack{j=1 \\ j \neq n}}^{\infty} \frac{H_{nj}}{j^2 - n^2} \sin(nx) \sin(j\tilde{t}) + \sum_{n=1}^{\infty} h_{nn} \sin(nx) \sin(n\tilde{t}),$$

where

$$H(x, \tilde{t}) \equiv 2\omega_1 \frac{\partial^2 \varphi_1(x, \tilde{t})}{\partial \tilde{t}^2} + 3\varphi_1(x, \tilde{t})\varphi_0^2(x, \tilde{t}) \equiv \sum_{n=1}^{\infty} \sum_{j=1}^{\infty} H_{nj} \sin(nx) \sin(j\tilde{t}).$$

It should be noted that all diagonal coefficients h_{nn} are arbitrary.

4 Conclusions

Using massless φ^4 theory as an example, we have shown that a uniform expansion of solutions for quasilinear Klein–Gordon equations can be constructed even in the principal resonance case. To construct the uniform expansion we have used the Poincaré–Lindstedt method and the nontrivial zero approximation: the function

$$\varphi_0(x, t) = A \left\{ \text{cn}(\alpha(x - \omega t), k) - \text{cn}(\alpha(x + \omega t), k) \right\},$$

with $k = 0.451075598811$ and $\alpha = 1.0576653982$.

Thus, using the Jacobi elliptic function **cn** instead of the trigonometric function **cos**, we have put the principal resonance to zero and constructed, with accuracy $\mathcal{O}(\varepsilon^3)$, the doubly periodic solution in the standing wave form:

$$\varphi(x, \omega t) = \varphi_0(x, \omega t) + \varepsilon \varphi_1(x, \omega t) + \varepsilon^2 \varphi_2(x, \omega t) + \mathcal{O}(\varepsilon^3),$$

with the frequency

$$\omega = 1 + \frac{\gamma^2}{64k^2} A^2 \varepsilon - \frac{\gamma^4}{8192k^4} A^4 \varepsilon^2 + \mathcal{O}(\varepsilon^3) = 1 + 1.0983600974 A^2 \varepsilon - 0.6031974518 A^4 \varepsilon^2 + \mathcal{O}(\varepsilon^3).$$

The question of constructing the uniform asymptotic series on the base of the obtained approximation remains open. The authors plan to investigate this question in future publications.

Acknowledgement

The authors are grateful to M. V. Tchitchikina, V. F. Edneral and P. K. Silaev for valuable discussions. S. Yu. Vernov would like to thank the Organizing Committee of the Fourth International IMACS Conference on Applications of Computer Algebra (ACA'98) for invitation to the Conference, hospitality and financial support.

This work has been supported by the Russian Foundation for Basic Research under Grants $\mathcal{N} \geq 96-15-96674$ and $\mathcal{N} \geq 98-02-27299$.

Appendix

This program have been written in **REDUCE 3.6**.

The program constructs system (7): $R_{jj} = 0$ and finds a particular solution of this system. The function $\varphi_0(x, \tilde{t})$ has been denoted as $\text{phi}(\text{xx}, \text{tt})$. This function is a solution of equation (5). We assume that the function $\text{phi}(\text{xx}, \text{tt})$ is real and includes only odd harmonics:

$$\text{phi}(\text{xx}, \text{tt}) = \sum_{j=1}^n a(2j-1) \sin((2j-1)\text{xx}) \sin((2j-1)\text{tt}).$$

The number of unknowns (parameter n) can be selected arbitrary. For $n = 10$ this program gives result on computer with 16 Mbytes operating memory, for $n = 20$ this program gives result on computer with 128 Mbytes operating memory.

Using the procedure "fourier" we expand equation (6) in the Fourier series and obtain the list of equations "listequa". The number of equations is $3n$. The unknowns $a(j)$ are a Fourier coefficients of some function, hence, we can assume that the sequence of the numbers $c(j) \equiv a(j)/a(1)$, satisfies the following condition: $\forall j > 1 : |c(j)| < 1$.

At the first step we assume that for $j > 3$, all $c(j) = 0$. From the first equation of "listequa" we find that C_{omega} is a polynomial in $c(3)$. We substitute this value of C_{omega} in the second equation of "listequa". As result we obtain that this equation is a cubic equation in $c(3)$. We solve this equation, using the standard procedure **SOLVE**. One of its solution must be real and we select this solution as value of $c(3)$.

At the second step we assume that $c(5)$ is arbitrary { clear $c(5)$ }. From the first equation of "listequa" we obtain that C_{omega} is a polynomial in $c(5)$, substitute this value of C_{omega} in the third equation of "listequa" and solve it, determining $c(5)$.

At the third step we substitute new values of $c(5)$ and C_{omega} in the second equation of "listequa", so we calculate the variable "equa". If "equa" is more than $\delta = 10^{-11}$, then we resolve the second equation of "listequa" and repeat the second step. Alternatively, if the obtained value of "equa" is less than or equal to $\delta = 10^{-11}$, then we assume that the three leading equations of "listequa" are solved and seek $c(7)$ from the fourth equation of "listequa" and so on. Once the coefficient $c(2n-1)$ has been found the program finishes to work.

```
out "rjj.res";
in fourier$
n:=10$
operator a,c,phi$
depend phi,xx,tt$
```

```
% Expand the unknown function phi(xx,tt), which is a solution of
% equation (5), and equation (6) in the Fourier series.
```

```
phi(xx,tt):=for j:=1:n sum a(2j-1)*sin((2j-1)*xx)*sin((2*j-1)*tt)$
equation1:=2*w1*df(phi(xx,tt),tt,2)+
```

```

                                fourier(fourier(phi(xx,tt)**3,xx),tt)$
listequa:={} $

% The frequency w_1 is proportional to a(1)^2.

w1:=C_omega*a(1)**2$
for j:=2:n do a(2*j-1):=c(2*j-1)*a(1)$
equation1:=equation1$

% Construct the system (7):  $R_{\{jj\}}(c)=0$ .

for k:=1:3*n do
    listequa:=append(listequa,
        {16*df(equation1,sin((2*k-1)*xx),
            sin((2*k-1)*tt))/a(1)**3});

% Find a real solution of this system.

on rounded$
C_omega:=0$

% C_omega must be connected variable, because I want to clear it
% without message:"WARNING...".

for k:=2:n do
<< for j:=k:n do c(2*j-1):=0;
    clear C_omega, c(2*k-1);
    firstequa:=first(listequa);
    C_omega:=C_omega-firstequa/df(firstequa,C_omega);
    equa:=part(listequa,k);

% All equations, which we solve using the procedure SOLVE, are
% cubic equations, hence, they have at least one real root. We assume
% that the absolute value of one of real roots is less than unit.
% We solve the equation and select this real solution
% as value of c(2*k-1).

    solve_equa:=solve(equa,c(2*k-1));
    while solve_equa neq {} do
<< c(2*k-1):=part(solve_equa,1,2);
        if( c(2*k-1)=sub(i=-i,c(2*k-1)) and abs(c(2*k-1))<1)
            then solve_equa:={}
            else solve_equa:=rest(solve_equa);
>>;
    test:=0;

```

```

while(test=0) do
<< test:=1;
  clear C_omega;
  firstequa:=first(listequa);
  C_omega:=C_omega-firstequa/df(firstequa,C_omega);
  for j:=2:k do
  <<
    equa:=part(listequa,j);
    if(abs(equa)>10**(-11)) then
    << clear c(2*j-1);
      test:=0;
      equa:=part(listequa,j);
      solve_equa:=solve(equa,c(2*j-1));
      while solve_equa neq {} do
        << c(2*j-1):=part(solve_equa,1,2);
          if(c(2*j-1)=sub(i=-i,c(2*j-1)) and abs(c(2*j-1))<1)
            then solve_equa:={}
            else solve_equa:=rest(solve_equa);
        >>;
      >>;
    >>;
  >>;
>>;

% Write the obtained result.

c(1):=1$
for j:=1:n do write "c(",2*j-1,"):= ", c(2*j-1);
for j:=2:n do write "c(",2*j-3,")/c(",2*j-1,"):= ", c(2*j-3)/c(2*j-1);
write "C_omega:= ", C_omega;
for j:=1:3*n do write "R(",j,",",j,"):= ",part(listequa,j);
quit;

```

The procedure "fourier" expands polynomials of $\sin(x)$ and $\cos(x)$ in Fourier series.

```

procedure fourier(FF,X);

% This procedure constructs Fourier-series expansions for polynomials
% of  $\sin(x)$  and  $\cos(x)$ .

begin
  scalar F;
  for all a, b such that df(a,x) neq 0 and df(b,x) neq 0
    let

```

```

cos(a)*cos(b)=(cos(a-b)+cos(a+b))/2,
sin(a)*sin(b)=(cos(a-b)-cos(a+b))/2,
sin(a)*cos(b)=(sin(a-b)+sin(a+b))/2,
sin(a)**2=(1-cos(2*a))/2,
cos(a)**2=(1+cos(2*a))/2;

F:=FF;
for all a, b such that df(a,x) neq 0 and df(b,x) neq 0
  clear
    cos(a)*cos(b),
    sin(a)*sin(b),
    sin(a)*cos(b),
    sin(a)**2,
    cos(a)**2;

return F;
end;
```

References

- [1] J. M. Coron, *Période Minimale Pour une Corde Vibrante de Longueur Infinie*, C. R. Acad. Sci. Paris Ser. A. **294** (1982) 127–129.
- [2] G. Lamb, *Elements of Soliton Theory*, Wiley, New York, **1980**.
- [3] P. Rabinowitz, *Free Vibrations for a Semilinear Wave Equation*, Commun. Pure Appl. Math. **31** (1978) 31–68.
- [4] H. Brezis, J. M. Coron, L. Nirenberg, *Free Vibrations for a Nonlinear Wave Equation and a Theorem of P. Rabinowitz*, Commun. Pure & Appl. Math. **33** (1980) 667–689.
- [5] H. Brezis, *Periodic Solutions of Nonlinear Vibrations Strings and Duality Principles*, Bulletin (New Series) of the AMS **8** (1983) 409–426.
- [6] G. Duffing, *Erzwungene Schwingungen bei Veränderlicher Eigenfrequenz und Ihre Technische Bedeutung*, Vieweg, Braunschweig, **1918**.
- [7] *Higher Transcendental Functions. Volume 3*. Edited by H. Bateman, A. Erdelyi. MC Graw-Hill Book Company, inc., New York, Toronto, London, **1955**.
- [8] S. Aubry, *A Unified Approach to the Interpretation of Displasive and Order-Disorder Systems II*, J. of Chem. Phys., **64** (1976) 3392–3402.
- [9] D. F. Kurdgelaidze, *Theory of the Nonlinear Field* $(\square - \lambda\varphi^2)\varphi = 0$, JETP (Rus. J. of Exper. & Theor. Phys.), **36** (1959) 842–849. {in Russian}
- [10] V. G. Bagrov, A. S. Vshivtsev, S. V. Ketov, *Additional Chapters of Mathematical Physics (Gauge Fields)*, Tomsk, Tomsk University Publishing House, **1990**. {in Russian}

- [11] O. A. Khrustalev, P. K. Silaev, *Doubly Periodic Solutions in Considerable Nonlinear 1-dimensional Field Model*, TMF (Rus. J. of Theor. & Math. Phys.), **117** (1998) 300–307. {in Russian}
- [12] H. Poincaré. *Oeuvres*, V. I–XI, Paris, Gauthier–Villars, **1934–1956**; *Les méthodes nouvelles de la mécanique céleste*, Paris, Gauthier–Villars, **1982**; *New Methods of Celestial Mechanics*, V. 1–3. NASA TTF–450, **1967**.
- [13] N. N. Bogoliubov, Yu. A. Mitropolskiy, *Asymptotic Methods in Theory of Nonlinear Oscillations*, Moscow, Nayka, **1974**. {in Russian}
- [14] A. H. Nayfeh, *Perturbation Methods*, John Wiley & Sons, New York, **1973**.
- [15] N. N. Moiseev, *Asymptotic Methods of Nonlinear Mechanics*, Moscow, Nauka, **1981**. {in Russian}
- [16] A. D. Bruno, *Local method in Nonlinear Differential Equations*, Springer Series in Soviet Mathematics, ISBN-3-540-18926-2, **1989**.
- [17] A. D. Bruno, *Power Geometry in Algebraic and Differential Equations*, Moscow, Nayka, **1998**. {in Russian}
- [18] A. C. Hearn, **REDUCE**. *User's Manual. Version 3.6*, RAND Publication CP78, **1995**.
- [19] V. F. Edneral, A. P. Kryukov, A. Ya. Rodionov, *The Language for Algebraic Computation REDUCE*, Moscow, MSU Publishing House, **1989**. {in Russian}
- [20] G. Rayna, **REDUCE**. *Software for Algebraic Computation*, New York: Springer, **1989**.
- [21] V. F. Edneral, O. A. Khrustalev, Proc. of the Int. Conf. on Computer Algebra and its Applications in Theoretical Physics. (USSR, Dubna, 1985), Dubna: JINR publ., (1985) 219–224. {in Russian}
- [22] V. F. Edneral, O. A. Khrustalev, *Program for Recasting ODE Systems in the Normal Form*, Programirovanie (Rus. J. of Programming) # **5** (1992) 73–80. {in Russian}
- [23] B. Buchberger, *Theoretical Basis for the Reduction of Polynomials to Canonical Forms.*, SIGSAM Bulletin **39** (1976) 19–29; *Some Properties of Gröbner–Bases for Polynomial Ideals*, SIGSAM Bulletin **40** (1976) 19–24.
- [24] J. Davenport, Y. Siret, E. Tournier, *Calcul Formel, Systemes et Algorithmes de Manipulations Algebriques*, Masson, Paris, New York, **1987**.
- [25] S. Yu. Vernov, O. A. Khrustalev, *Approximate Doubly Periodic Solutions in $(1+1)$ -dimensional φ^4 Theory*, TMF (Rus. J. of Theor. Math. Phys.) **116** (1998) 182–192 {in Russian}; 881–889 {in English}.